

RECOMMENDED CLINICAL PREVENTIVE SERVICES FOR ADULT MEN

Clinical preventive services for adults, based on the AAFP Summary of Recommendations for Clinical Preventive Services



	18	25	30	35	40	45	50	55	60	65	70	75
ACCIDENTAL INJURY	Ⓡ Counsel as appropriate for age											
ALCOHOL MISUSE	Ⓡ Screen and counsel behavior to reduce misuse											
AORTIC ANEURYSM, ABDOMINAL	Ⓡ One-time screening by ultrasonography ¹											
COLORECTAL CANCER	Ⓢ Screen for colorectal cancer											
CORONARY HEART DISEASE	Ⓢ Counsel adults at increased risk regarding benefits and risks of aspirin prophylaxis											
DEPRESSION	Ⓡ Screen for depression											
HEARING DIFFICULTIES	Ⓡ Screen and counsel											
HYPERTENSION	Ⓢ Screen for high blood pressure											
LIPID DISORDERS	Ⓢ Screen with fasting lipid profile or nonfasting total and HDL cholesterol											
OBESITY	Ⓡ Screen for obesity by measuring height and weight periodically											
	Ⓢ Intensive counseling and behavioral interventions to promote sustained weight loss for obese adults ²											
PHYSICAL ACTIVITY	Ⓢ Recognize that physical activity is desirable and provide advice accordingly											
SECONDHAND SMOKE	Ⓢ Counsel parents who smoke regarding harmful effects of smoking on children's health											
STDs	Ⓡ Counsel regarding the risks for STDs and how to prevent them											
TOBACCO USE	Ⓢ Screen for tobacco use and provide tobacco cessation interventions as appropriate											
VIOLENCE, FAMILY AND PARTNER	Ⓢ Be alert to physical and behavioral signs and symptoms associated with abuse or neglect											
VISUAL DIFFICULTIES	Ⓡ Screen with Snellen acuity											

UPPER AGE LIMITS SHOULD BE INDIVIDUALIZED FOR EACH PATIENT

Ⓢ Strongly Recommend: Good quality evidence exists which demonstrates substantial net benefit over harm; the intervention is perceived to be cost effective and acceptable to nearly all patients.

Ⓡ Recommend: Although evidence exists which demonstrates net benefit, either the benefit is only moderate in magnitude or the evidence supporting a substantial benefit is only fair. The intervention is perceived to be cost effective and acceptable to most patients.

Ⓢ Healthy Behavior: Healthy Behavior is identified as desirable but the effectiveness of physician's advice and counseling is uncertain.

AAFP Age Charts for Clinical Preventive Services are based on the AAFP Summary of Recommendations for Clinical Preventive Services. These charts include only positive recommendations, detailed language and further information, consult the Recommendations. For immunization information, you may also consult the Adult Immunization Schedule. These age charts are provided only as an assistance for physicians making clinical decisions regarding the care of their patients. They cannot substitute for the individual judgment brought to each clinical situation by the patient's family physician. Based on AAFP Policy Action November 1996; Revision 6.1, April 2006.

Copyright © 2006 American Academy of Family Physicians. Permission is granted to reproduce this material for nonprofit educational uses. Written permission is required for all other uses, including electronic uses.

1. In men who have ever smoked

2. Intensive counseling involves more than 1 session per month for at least 3 months